

A Helpful Diagnostic Tool

RECOMMENDATIONS

- All women should be assessed for a history of depression in the prenatal period (including evaluation of a history of prepartum psychosis or family history of affective disorders).
- Edinburgh Postnatal Depression Screen should be offered postpartum as *part of* a screening program for postpartum depression. The diagnosis requires clinical evaluation in addition to this screening.
- The Edinburgh Postnatal Depression Screen should be given at six weeks and three months following delivery.
- Women at high risk should have prenatal education and more frequent postpartum visits. ^{1,2}

Edinburgh Postnatal Depression Screen				
<i>*Reproduced with permission from the British Journal of Psychiatry, June 1987. Vol. 150; J. L. Cox; J. M. Holden and R. Sagovsky</i>				
Name:				
Address:				
Baby's Age:				
As you have recently had a baby, we would like to know how you are feeling. Please check the answer which comes closest to how you have felt in the PAST 7 DAYS, not just how you feel today,				
	Yes very often	Sometimes	Hardly ever	No, Not at all
1. I have been able to laugh and see the funny side of things.	0	1	2	3
2. I have looked forward with enjoyment to things.	0	1	2	3
3. I have blamed myself unnecessarily when things went wrong.	3	2	1	0
4. I have felt worried or anxious without a very good reason.	3	2	1	0
5. I have felt scared or panicky without a good reason.	3	2	1	0
6. I have been feeling overwhelmed	3	2	1	0
7. I have have had difficulty sleeping even when the baby is asleep	3	2	1	0
8. I have felt sad or miserable.	3	2	1	0
9. I have been so unhappy that I have been crying, or fighting to keep from crying.	3	2	1	0

10. The thought of harming either myself or my baby has occurred to me.	3	2	1	0
TOTAL				
Scoring				
0-8 points	Low probability of Depression			
8-12 points	Most likely dealing with life with a new baby or a case of post partum blues, also known as “baby blues”			
13-14 points	Signs leading to the possibility of post partum depression; take preventive measures			
15+ points	High probability of experiencing clinical post partum depression			

Any reference in this material to other organizations or companies, including their Internet Web sites, is not an endorsement or warranty of services, information or products provided by those organizations or companies.

This information has been provided to you by CIGNA Behavioral Health. It was collaboratively created by CIGNA Behavioral Health and CIGNA HealthCare and published originally as part of the CIGNA HealthCare Healthy Babies program. Nothing in this information should be construed as a specific recommendation for medical care or treatment. The information in this document is not intended to be a substitute for your professional judgment or experience, but is designed to provide you with additional tools and information that support your efforts and make it easier for you to deliver quality care.

"CIGNA HealthCare" refers to various operating subsidiaries of CIGNA Corporation. Products and services are provided by these subsidiaries and not by CIGNA Corporation. These subsidiaries include Connecticut General Life Insurance Company, Tel-Drug, Inc. and its affiliates, CIGNA Behavioral Health, Inc., Intracorp, and HMO or service company subsidiaries of CIGNA Health Corporation and CIGNA Dental Health, Inc.

¹ Scottish Intercollegiate Guideline Network, Royal College of Physicians, Edinburgh, 2002.

² Cox, J.L., Holden, J.M. and Sagovsky, R., *British Journal of Psychiatry*, June 1987. Vol. 150.